

TSKgel[®] SP-STAT and CM-STAT Columns

INTRODUCTION

TSKgel SP-STAT and TSKgel CM-STAT cation exchange columns allow fast equilibration and analysis, as well as isolation, of complex biomolecules. Both TSKgel columns are packed with 7 or 10 μm mono-disperse, non-porous resin particles of which the surface consists of an open access network of multi-layered cation exchange groups (see Figure 1). The innovative bonding chemistry, combined with a relatively large particle size, result in a respectable loading capacity and a low operating pressure, attributes not found in traditional mono-disperse, non-porous resins.


PRODUCT HIGHLIGHTS

- Very efficient chromatography for high as well as low MW solutes made possible by novel bonding chemistry and the absence of micro-pores
- High speed and high resolution analysis of biomolecules
- Higher adsorption capacities and lower pressures compared to competitive non-porous columns
- 7 or 10 μm particles for SP and CM chemistries

APPLICATIONS

FAST SEPARATIONS

The fast separation of protein standards was investigated using short cation exchange columns (see Figure 2). A TSKgel SP-STAT column shows superior resolution, better peak shape, and a shorter analysis time (< 60 seconds) compared to a competitive monolithic SP-type column.




➤ Figure 1

REACTION MONITORING

A sample of β -lactoglobulin (5 mg/mL) was reacted with polyethylene glycol (5 kDa) in a pH 6.5 phosphate buffer. The formation of PEGylated protein reaction products was monitored in 5 minute intervals on a 3.5 cm TSKgel SP-STAT column. As demonstrated in Figure 3, peak areas of mono-, di-, and tri-PEGylated β -lactoglobulin increased with reaction time, while the area of unreacted β -lactoglobulin declined.

ANTIBODY ANALYSIS

The analysis profiles for five antibodies separated on a TSKgel CM-STAT column were compared with the profiles obtained on a competitive WCX column (Figure 4). Similar or higher resolution profiles were obtained on TSKgel CM-STAT in approximately half the time.



➤ Figure 2

Column: A: TSKgel SP-STAT, 10 μm , 3.0 mm ID x 3.5 cm L; B: Competitor column 4.6 mm ID x 5.0 cm L
 Eluent: A: 20 mmol/l sodium acetate (pH 5.0); B: 1.0 mol/l NaCl in buffer A (pH 5.0) for column A; 1.5 mol/l NaCl in buffer A (pH 5.0) for column B;
 Gradient: 0% B (0 min), 100% B (1 min);
 Flow rate: A: 2.0 ml/min; B: 4.73 ml/min; Detection: UV @ 280 nm;
 Samples: 1. α -chymotrypsinogen A; 2. cytochrome C; 3. lysozyme




Figure 3

Column: TSKgel SP-STAT, 10 μ m, 3.0 mm ID x 3.5 cm L; Eluent: A: 20 mmol/l sodium acetate (pH 5.0); B: 1.0 mol/l NaCl in buffer A (pH 5.0); Gradient: 0% B (0 min), 100% B (2 min); Flow rate: 2.0 ml/min; Detection: UV @ 280 nm; Samples: PEGylated β -lactoglobulin




Figure 4

Column: A: TSKgel CM-STAT, 7 μ m, 4.6 mm ID x 10 cm L; B: Competitor WCX, 10 μ m, 4.0 mm ID x 25 cm L; Eluent: A: 20 mmol/L MES (pH 6.0); B: 20 mmol/L MES + 0.5 mol/L NaCl (pH 6.0) Gradient: A: 10% B (0 min), 30% B (15 min), 100% B (15 min), 100% B (17 min), 10% B (17 min), 10% B (21 min); B: 10% B (0 min), 30% B (30 min), 100% B (30 min), 100% B (32 min), 10% B (32 min), 10% B (36 min) Flow rate: A: 1.0 mL/min B: 2.0 mL/min; Temp.: Ambient; Detection: UV @ 280 nm; Inj. Vol.: 20 μ L; Sample: monoclonal antibodies (mAb A through E)

Ordering information

TSKgel STAT COLUMNS

Part-No	Description	Matrix	Housing	Dimensions
0021963	TSKgel SP-STAT, 10 μ m	Polymer	Stainless steel	3.0 mm ID x 3.5 cm L
0021964	TSKgel SP-STAT, 7 μ m	Polymer	Stainless steel	4.6 mm ID x 10 cm L
0021965	TSKgel CM-STAT, 10 μ m	Polymer	Stainless steel	3.0 mm ID x 3.5 cm L
0021966	TSKgel CM-STAT, 7 μ m	Polymer	Stainless steel	4.6 mm ID x 10 cm L